

A note on the normal largest gap between prime factors

Gérald Tenenbaum

Abstract. Let $\{p_j(n)\}_{j=1}^{\omega(n)}$ denote the increasing sequence of distinct prime factors of an integer n . We provide details for the proof of a statement of Erdős implying that, for any function $\xi(n)$ tending to infinity with n , we have

$$f(n) := \max_{1 \leq j < \omega(n)} \log \left(\frac{\log p_{j+1}(n)}{\log p_j(n)} \right) = \log_3 n + O(\xi(n))$$

for almost all integers n .

Keywords: Distribution of prime factors, normal order, largest gap.

2010 Mathematics Subject Classification: 11N25, 11N35.

In private correspondence [4], E. Sofos asked for normal upper and lower bounds for the arithmetic function

$$f(n) := \max_{1 \leq j < \omega(n)} \log \left(\frac{\log p_{j+1}(n)}{\log p_j(n)} \right),$$

where $\{p_j(n)\}_{j=1}^{\omega(n)}$ denotes the increasing sequence of distinct prime factors of an integer n . The answer to this question is actually provided by a statement of Erdős in [1], where it is asserted without proof that, for any $c > 0$, the density of the set of those integers n such that $e^{f(n)} > c \log_2 n$ is $1 - e^{-1/c}$. Here and in the sequel \log_k denotes the k -fold iterated logarithm. Indeed, given any function $\xi(n)$ tending to infinity and letting c tend sufficiently slowly to zero, we infer that, for almost all n , we have $f(n) > \log_3 n - \xi(n)$, while by letting c tend slowly to infinity we obtain that $f(n) \leq \log_3 n + \xi(n)$ also holds for a set of asymptotic density 1. Interesting related results appear in Erdős' articles [2] and [3].

In this short note, we provide the not so obvious details of the proof, in the spirit of the paper [1].

Theorem 1 (Erdős [1]). *Let $c > 0$. Then the inequality $f(n) > \log_3 n - \log(1/c)$ holds on a set of integers n of asymptotic density $1 - e^{-1/c}$. In particular, given any function $\xi(n) \rightarrow \infty$, we have*

$$\log_3 n - \xi(n) \leq f(n) \leq \log_3 n + \xi(n)$$

for almost all integers n .

Proof. Let x be a large parameter, write $Z := c \log_2 x$, and let $\chi_p(n)$ denote the indicator function of the set of those integers $n \leq x$ that are divisible by the prime p but by no prime q such that $p < q \leq p^Z$. For squarefree m , put $\chi_m(n) := \prod_{p|m} \chi_p(n)$. Taking into account the slow growth of $\log_2 x$, it is sufficient to show that

$$(1) \quad N(x) := |\{n \leq x : p \mid n, p \leq x^{1/Z} \Rightarrow \chi_p(n) = 0\}| \sim e^{-1/c} x \quad (x \rightarrow \infty).$$

Indeed, $n \leq x$ is counted by $N(x)$ if, and only if, $p_{j+1}(n) \leq p_j(n)^Z$ whenever $1 \leq j < \omega(n)$. Let $P^+(n)$ (resp. $P^-(n)$) denote the largest (resp. the smallest) prime factor of an integer n , with the convention that $P^+(1) = 1$ (resp. $P^-(1) = \infty$), and let μ designate the Möbius function. By the inclusion–exclusion principle (see, e.g., [6], p. 39) or the Möbius inversion formula, we thus have, for $x \geq 1$,

$$(2) \quad N(x) = \sum_{\substack{m \leq x \\ m \in \mathcal{M}}} \mu(m) \sum_{n \leq x} \chi_m(n),$$

where \mathcal{M} denotes the set of all squarefree integers $m \leq x$ such that $P^+(m) \leq x^{1/Z}$ and $p_{j+1}(m) > p_j(m)^Z$ whenever $j \in [1, \omega(m)[$. Moreover, as is familiar in Brun's sieve method, we obtain a lower or an upper bound for the left-hand side by restricting the outer sum to integers m with number of prime factors at most equal to an odd or an even bound.

Let $k \geq 1$ be fixed. For each $m \in \mathcal{M}$, define $P_m := \prod_{p|m} \prod_{p < q \leq p^z} q$. Then

$$(3) \quad N_k(x) := \sum_{\substack{m \in \mathcal{M} \\ \omega(m)=k}} \sum_{n \leq x} \chi_m(n) = \sum_{\substack{m \in \mathcal{M} \\ \omega(m)=k}} \sum_{\substack{\nu \leq x/m \\ (\nu, P_m)=1}} 1.$$

From a standard sieve result (see, e.g., the lemma in [5]), we infer that the inner sum is

$$(4) \quad \sim \frac{x\varphi(P_m)}{mP_m} = \frac{x}{mZ^k} \left\{ 1 + O\left(\frac{1}{\log P^-(m)}\right) + o(1) \right\}$$

provided $P^+(m) \leq x^{o(1/Z)}$. The contribution of the remaining integers m will be treated as an error term. Indeed, in this last case, we may plainly assume that $x^{\varepsilon_x} < P^+(m)^Z \leq x$, with ε_x tending to 0 arbitrarily slowly. Then the inner sum is classically (see, e.g., [6], Exercise 85)

$$\ll \frac{x\varphi(P_m)}{mP_m} \ll \frac{x}{mZ^k}$$

and the corresponding contribution to $N_k(x)$ is

$$\frac{x}{Z^k} \sum_{x^{\varepsilon_x/Z} < p \leq x^{1/Z}} \frac{1}{p} \sum_{\substack{P^+(h) \leq x \\ \omega(h)=k-1}} \frac{1}{h} \ll \frac{x \log(1/\varepsilon_x)}{\log_2 x} = o(x).$$

Inserting (4) back into (3) and summing over m , we obtain, for $k \geq 0$,

$$N_k(x) = \frac{x}{Z^k} \sum_{\substack{p_k \leq x^{1/Z} \\ p_j \leq p_{j+1}^{1/Z} (1 \leq j < k)}} \frac{1 + O(1/\log p_1) + o(1)}{p_1 \cdots p_k} \sim \frac{x(\log_2 x)^k}{k!Z^k} \sim \frac{x}{c^k k!}.$$

Thus, for arbitrary $\ell \geq 1$, we have

$$x \sum_{0 \leq k \leq 2\ell+1} \frac{(-1)^k}{c^k k!} + o(x) \leq N(x) \leq x \sum_{0 \leq k \leq 2\ell} \frac{(-1)^k}{c^k k!} + o(x),$$

and the required result follows by selecting ℓ arbitrarily large. \square

Acknowledgements. The author wishes to express warm thanks to Régis de la Bretèche for valuable discussions on this matter.

References

- [1] P. Erdős, Some remarks on prime factors of integers. *Canad. J. Math.* **11** (1959), 161–167.
- [2] P. Erdős, On some properties of prime factors of integers, *Nagoya Math. J.* **27** (1966), 617–623.
- [3] P. Erdős, On the distribution of prime divisors, *Aequationes Math.* **2** (1969), 177–183.
- [4] E. Sofos, private e-mail message, August 30, 2018.
- [5] G. Tenenbaum, Crible les entiers sans grand facteur premier, in: R. C. Vaughan (ed.), *Theory and applications of numbers without large prime factors*, *Phil. Trans. R. Soc. Lond. A* **345** (1993), 377–384.
- [6] G. Tenenbaum, *Introduction to analytic and probabilistic number theory*, 3rd ed., Graduate Studies in Mathematics 163, Amer. Math. Soc. 2015.

Gérald Tenenbaum
 Institut Élie Cartan
 Université de Lorraine
 BP 70239
 54506 Vandœuvre-lès-Nancy Cedex
 France
 gerald.tenenbaum@univ-lorraine.fr